

ENBREL[®] PATIENT PASSPORT

Important safety information
on your treatment
with Enbrel[®] (etanercept)

Enbrel[®]
Etanerceptum

TABLE OF CONTENTS

| | |
|---|----|
| What you should be aware of before use..... | 3 |
| Tell your doctor that you are being treated with Enbrel®..... | 4 |
| Infections..... | 6 |
| Your medication..... | 8 |
| Batch number record..... | 10 |
| Treating physician..... | 14 |
| Information on the patient..... | 15 |

WHAT YOU SHOULD BE AWARE OF BEFORE USE

This patient passport contains important safety information that you need to be aware of before and during treatment with Enbrel®. If you do not understand this information, please ask your doctor to explain it to you.

- Show this card to any doctor involved in your treatment.
- See the Enbrel® package leaflet for more information.
- Keep this patient passport with you for 2 months after your last dose of Enbrel® as side effects may still occur after the final dose.

TELL YOUR DOCTOR THAT YOU ARE BEING TREATED WITH ENBREL®

- prior to major surgery (and at the dentist)
- in the presence of infections, persistent fever, inflammation of the throat, bruises, bleeding or pallor, and that tuberculosis screening must be conducted
- in the presence of allergic reactions, such as tightness in the chest, wheezing, drowsiness or skin rash
- if there are signs of heart failure (cardiac insufficiency), such as shortness of breath at night or during exercise, coughing at night, swelling of the ankles, feeling of pressure in the neck/abdominal region, blue discolouration of the nails/lips

- if you plan to get pregnant, are pregnant or breast-feeding. A waiting period of 16 weeks after the mother's last dose of Enbrel® is recommended before the administration of live vaccines are given to infants that were exposed to Enbrel® *in utero*, unless serum Enbrel® concentrations can no longer be detected in the infant or the benefits of the vaccination significantly outweigh the theoretical risk of administering live vaccines to infants "Either breastfeeding should be paused during treatment, or treatment should be paused during breastfeeding".
- prior to vaccinations and before you take other medication, including over-the-counter medication

Important note: This list is not exhaustive.

Please consult the Information for Patients and/or ask your doctor/pharmacist or a nurse.

INFECTIONS

**Enbrel® may increase your risk of infection.
These infections may also be serious.**

- Do not take Enbrel® if you have an infection. Please ask your doctor if you are unsure.
- You must get medical treatment immediately if you develop any symptoms that may point to an infection - e.g. fever, persistent cough, weight loss or apathy.
- You should be screened for tuberculosis (TB) before you start treatment. Please ask your doctor to record the data and results of your last screening for TB below:

Test

Date

Result

YOUR MEDICATION

Please ask your doctor to list your other medication that might increase your risk of infection.

BATCH NUMBER RECORD

It is important that you record the batch numbers printed on the pack for all forms of treatment with Enbrel®:

I am using an:

- Enbrel® pre-filled syringe
- Enbrel® MYCLIC® pre-filled pen
- Enbrel® lyophilisate

Date of use

Batch/Lot number

BATCH NUMBER RECORD

Date of use

Batch/Lot number

Date of use

Batch/Lot number

TREATING PHYSICIAN

Name:

Address:

Phone no.:

Fax no.:

INFORMATION ON THE PATIENT

Name:

Address:

Phone no.:

Date of birth:



Pfizer AG
Schärenmoosstrasse 99
Postfach
8052 Zürich

PP-ENB-CHE-0661 Dez 2023